

---

**Accelerated Article Preview**

---

# Geological Net Zero and the need for disaggregated accounting for carbon sinks

---

Received: 2 March 2024

---

Accepted: 31 October 2024

---

Accelerated Article Preview

---

Cite this article as: Allen, M. R. et al. Geological Net Zero and the need for disaggregated accounting for carbon sinks. *Nature* <https://doi.org/10.1038/s41586-024-08326-8> (2024)

Myles R. Allen, David J. Frame, Pierre Friedlingstein, Nathan P. Gillett, Giacomo Grassi, Jonathan M. Gregory, William Hare, Jo House, Chris Huntingford, Stuart Jenkins, Chris D. Jones, Reto Knutti, Jason A. Lowe, H. Damon Matthews, Malte Meinshausen, Nicolai Meinshausen, Glen P. Peters, Gian-Kasper Plattner, Sarah Raper, Joeri Rogelj, Peter A. Stott, Susan Solomon, Thomas F. Stocker, Andrew J. Weaver & Kirsten Zickfeld

---

This is a PDF file of a peer-reviewed paper that has been accepted for publication. Although unedited, the content has been subjected to preliminary formatting. Nature is providing this early version of the typeset paper as a service to our authors and readers. The text and figures will undergo copyediting and a proof review before the paper is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers apply.

# Geological Net Zero and the need for disaggregated accounting for carbon sinks

Myles R. Allen<sup>1,2,\*</sup>, David J. Frame<sup>3</sup>, Pierre Friedlingstein<sup>4</sup>, Nathan P. Gillett<sup>5</sup>, Giacomo Grassi<sup>6</sup>, Jonathan M. Gregory<sup>7,11</sup>, William Hare<sup>8</sup>, Jo House<sup>9</sup>, Chris Huntingford<sup>10</sup>, Stuart Jenkins<sup>2</sup>, Chris D. Jones<sup>11</sup>, Reto Knutti<sup>12</sup>, Jason A. Lowe<sup>13</sup>, H. Damon Matthews<sup>14</sup>, Malte Meinshausen<sup>15</sup>, Nicolai Meinshausen<sup>16</sup>, Glen P. Peters<sup>17</sup>, Gian-Kasper Plattner<sup>18</sup>, Sarah Raper<sup>19</sup>, Joeri Rogelj<sup>20</sup>, Peter A. Stott<sup>11,21</sup>, Susan Solomon<sup>22</sup>, Thomas F. Stocker<sup>23</sup>, Andrew J. Weaver<sup>24</sup>, and Kirsten Zickfeld<sup>25</sup>

\* Corresponding author e-mail: [myles.allen@physics.ox.ac.uk](mailto:myles.allen@physics.ox.ac.uk)

## Author affiliations:

1. Atmospheric, Oceanic and Planetary Physics, Department of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, UK
2. Oxford Net Zero, Environmental Change Institute, School of Geog. & the Environment, University of Oxford, S Parks Rd, Oxford OX1 3QY, UK
3. School of Physical and Chemical Sciences, University of Canterbury, PB 4800, Christchurch 8140, New Zealand
4. Faculty of Environment, Science and Economy, University of Exeter, Exeter EX4 4QF, UK
5. Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, BC, V8N 1V8, Canada
6. European Commission, Joint Research Centre (JRC), 21027 Ispra, Italy
7. National Centre for Atmospheric Science, Department of Meteorology, University of Reading, Reading RG6 6ET, UK
8. Climate Analytics, Ritterstraße 3, Berlin 10969, Germany
9. School of Geographical Sciences, University of Bristol, University Road, Clifton, Bristol, BS8 1SS, UK
10. UK Centre for Ecology and Hydrology, Wallingford, OX10 8BB, UK
11. The Met Office, Fitzroy Road, Exeter, EX1 3PB, UK
12. Institute for Atmospheric and Climate Science, ETH Zurich, 8092, Zürich, Switzerland
13. Priestley Centre for Climate Futures, University of Leeds, Priestley Building, Woodhouse, Leeds LS2 9JT, UK
14. Department of Geography, Planning & Environment, Concordia University, Montréal, QC, Canada
15. School of Geography, Earth and Atmospheric Sciences, University of Melbourne, Melbourne, Victoria, Australia
16. Seminar for Statistics, Department of Mathematics, ETH Zurich, 8092 Zurich, Switzerland
17. CICERO Center for International Climate Research, Oslo 0349, Norway
18. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, 8903 Birmensdorf, Switzerland
19. Manchester Metropolitan University, Ormond Building, Lower Ormond Street, Manchester M15 6BX, UK
20. Centre for Environmental Policy and Grantham Institute, Imperial College London, Exhibition Road, London SW7 2AZ, UK
21. Department of Mathematics, University of Exeter, Exeter, EX4 4QF, UK
22. Department of Earth, Atmospheric, and Planetary Sciences, MIT, 77 Massachusetts Ave., Cambridge, MA 02139, USA
23. Oeschger Centre for Climate Change Research, University of Bern, Hochschulstrasse 4, 3012 Bern, Switzerland
24. School of Earth and Ocean Sciences, University of Victoria, PO Box 1700, Victoria, B.C., V8W 2Y2 Canada
25. Department of Geography, Simon Fraser University, 8888 University Drive, Burnaby, B.C., V5A 1S6, Canada

53 **Preface:** Achieving net zero global emissions of carbon dioxide (CO<sub>2</sub>), with declining emissions of  
54 other greenhouse gases, is widely expected to halt global warming. CO<sub>2</sub> emissions will continue to  
55 drive warming until fully balanced by active anthropogenic CO<sub>2</sub> removals. For practical reasons,  
56 however, many greenhouse gas accounting systems allow some “passive” CO<sub>2</sub> uptake, such as  
57 enhanced vegetation growth due to CO<sub>2</sub> fertilisation, to be included as removals in the definition of  
58 net anthropogenic emissions. By including passive CO<sub>2</sub> uptake, nominal net zero emissions would not  
59 halt global warming, undermining the Paris Agreement. Here we discuss measures addressing this  
60 problem, to ensure residual fossil fuel use does not cause further global warming: land management  
61 categories should be disaggregated in emissions reporting and targets to better separate the role of  
62 passive CO<sub>2</sub> uptake; where possible, claimed removals should be additional to passive uptake; and  
63 targets should acknowledge the need for Geological Net Zero, meaning one tonne of CO<sub>2</sub> permanently  
64 restored to the solid Earth for every tonne still generated from fossil sources. We also argue that  
65 scientific understanding of net zero provides a basis for allocating responsibility for the protection of  
66 passive carbon sinks during and after the transition to Geological Net Zero.

67  
68 **The Problem:** The UAE Consensus<sup>1</sup>, agreed at the COP28 climate conference, called on Parties “to  
69 achieve net zero by 2050 in keeping with the science” without specifying precisely to what net zero  
70 refers.<sup>2</sup> The concept dates back to a series of papers<sup>3–8</sup> in 2009 that established the cumulative impact  
71 of anthropogenic carbon dioxide (CO<sub>2</sub>) emissions on global temperatures, and the need to reduce net  
72 CO<sub>2</sub> emissions to zero to halt global warming. This was affirmed<sup>9</sup> in the Intergovernmental Panel on  
73 Climate Change (IPCC)’s 5<sup>th</sup> Assessment Report (AR5) which informed Article 4.1 of the Paris  
74 Agreement: “In order to achieve the long-term temperature goal set out in Article 2 (“Holding the  
75 increase in the global average temperature to well below 2°C above pre-industrial levels and pursuing  
76 efforts to limit the temperature increase to 1.5°C”), Parties aim ... to achieve a balance between  
77 anthropogenic emissions by sources and removals by sinks of greenhouse gases in the second half of  
78 this century”. This wording, the foundation of subsequent national and corporate<sup>10</sup> net zero pledges,  
79 makes clear that the purpose of “balance” is to limit global warming. The IPCC’s Special Report on  
80 1.5°C (SR1.5)<sup>11</sup> stated what this entails: “Reaching and sustaining net-zero global anthropogenic CO<sub>2</sub>  
81 emissions and declining net non-CO<sub>2</sub> radiative forcing would halt anthropogenic global warming on  
82 multi-decadal timescales (*high confidence*)”, reaffirmed by subsequent research<sup>12,13</sup> and the IPCC 6<sup>th</sup>  
83 Assessment (AR6).<sup>14–16</sup>

84  
85 It is, however, increasingly clear that many current interpretations of net zero CO<sub>2</sub> emissions, if  
86 applied globally, are not consistent with the goal of halting the rise in global temperatures.<sup>17–19</sup> The  
87 problem is ambiguity in the definition of anthropogenic CO<sub>2</sub> removals (called “removals” for brevity  
88 hereon). The definition of removal used in IPCC Scientific Assessments<sup>20</sup> explicitly “excludes natural  
89 CO<sub>2</sub> uptake not directly caused by human activities” (here we use IPCC Scientific Assessment  
90 definitions<sup>20</sup> unless otherwise specified). Yet methods used by many greenhouse gas reporting  
91 systems, including those informed by the IPCC guidelines for national greenhouse gas inventories  
92 (NGHGs),<sup>21</sup> implicitly allow indirect or passive uptake (so-called because it is occurring as a  
93 consequence of past emissions and not as a result of active ongoing human intervention) to be classed  
94 as a removal if it takes place on “managed land”.<sup>22–24</sup> The concept of managed land was originally  
95 introduced, in part, because differentiating between active land-based removal of atmospheric CO<sub>2</sub>  
96 and passive CO<sub>2</sub> uptake<sup>25</sup> requires modelling a counterfactual i.e. what would have happened if the  
97 action leading to a claimed land-based removal had not occurred? This cannot be inferred from  
98 observations alone. Model-based approaches<sup>23</sup> allow a global mapping between different removal  
99 classification systems, but ambiguities remain, such as the classification of ongoing regrowth  
100 following reforestation. As pressure to reduce net emissions rises, more land may be deemed  
101 managed, reclassifying passive uptake as active removal. Already, not all claimed land-based CO<sub>2</sub>  
102 emission reductions<sup>26</sup> and removals<sup>27</sup> are verifiably additional to what would have occurred without  
103 any active human intervention. These problems are compounded by the risk of terrestrial carbon  
104 stocks being re-released through Earth system feedbacks. Similar problems may arise in the future  
105 with an increased focus on “blue carbon”<sup>31</sup> uptake by the oceans.

106

107 Hence, under the Global Stocktake,<sup>1</sup> pathways to net-zero are determined by models that use a narrow  
108 definition of CO<sub>2</sub> removals, excluding<sup>20</sup> all passive uptake, yet countries<sup>32</sup> and corporations<sup>10,27</sup>  
109 typically assess their progress using the broader NGHGI definition, which includes some passive  
110 uptake. If the definition of anthropogenic removals includes passive uptake then nominal “net zero”  
111 CO<sub>2</sub> emissions could fail to halt global warming in time to deliver the goals of the Paris Agreement.  
112

113 **Scientific context:** CO<sub>2</sub>-induced warming  $\Delta T_{\text{CO}_2}$  over a multi-decade time-interval  $\Delta t$  (such as 2025-  
114 2050, or 2050-2100) is, to a good approximation, given by<sup>18</sup>

$$\Delta T_{\text{CO}_2} = \kappa_E [E_{\text{GEO}} + E_{\text{LUC}} + (\rho_F - \rho_E)G] \Delta t . \quad (1)$$

118 The variables, affected by policy, are  $E_{\text{GEO}}$ , the average global net rate of geological-origin CO<sub>2</sub>  
119 emissions over that time-interval (total CO<sub>2</sub> produced from fossil fuels and industrial processes minus  
120 CO<sub>2</sub> captured at source or recaptured from the atmosphere and committed to permanent geological  
121 storage, in billions of tonnes per year);  $E_{\text{LUC}}$ , the net biogenic CO<sub>2</sub> emissions that result from ongoing  
122 direct anthropogenic land-use change (e.g., active deforestation, afforestation, reforestation and  
123 ecosystem restoration, including coastal habitats<sup>33,34</sup>), but not including passive (indirect) uptake  
124 driven by past emissions<sup>35</sup> (including CO<sub>2</sub> fertilisation of existing forests as well as temperature,  
125 precipitation, and growing season effects); and  $G$ , cumulative net CO<sub>2</sub> emissions that have resulted  
126 directly from all human activities from pre-industrial times up to the mid-point of the time-interval in  
127 question, in billions of tonnes. Total human-induced warming comprises  $\Delta T_{\text{CO}_2}$  plus non-CO<sub>2</sub>  
128 warming (see Methods).  
129

130 The coefficients, not affected by policy, are  $\kappa_E$ , the Transient Climate Response to Emissions  
131 (TCRE)<sup>8,20</sup>;  $\rho_F$ , the fractional Rate of Adjustment to Constant Forcing (RACF)<sup>18,36,37</sup>; and  $\rho_E$ , the  
132 Slow Carbon-cycle Adjustment Rate<sup>18</sup> or the fractional rate of CO<sub>2</sub> radiative forcing<sup>20</sup> decline under  
133 zero emissions.<sup>38,39</sup> Both rates are approximately 0.3% per year.<sup>16,40</sup> Equation 1 reproduces, within  
134 uncertainties due to internal climate variability, the response of coupled climate-carbon-cycle models  
135 to a broad range of emissions scenarios up to the time of peak warming.<sup>13</sup> Limiting CO<sub>2</sub>-induced  
136 warming, or reducing  $\Delta T_{\text{CO}_2}$  to zero, is necessary to halt total greenhouse-gas-induced global warming  
137 on multi-decadal timescales, while reductions in other greenhouse gas emissions are also required to  
138 meet Paris temperature goals. Henceforth, net zero refers to net zero CO<sub>2</sub> emissions unless specified  
139 otherwise.  
140

141 The first insight of the 2009 papers was that  $\kappa_E$  is largely time- and scenario-independent,<sup>9,15,41-43</sup> so  
142 that cumulative CO<sub>2</sub> emissions since pre-industrial times determine the level of CO<sub>2</sub>-induced  
143 warming.<sup>44</sup> The second was that  $\rho_E \approx \rho_F$ , so the difference between them, or Rate of Adjustment to  
144 Zero Emissions,<sup>13,18</sup> is approximately zero.<sup>12</sup> This cancellation means that no substantial further CO<sub>2</sub>-  
145 induced warming or cooling of the climate system will occur as long as  $E_{\text{GEO}} + E_{\text{LUC}} = 0$ . These two  
146 findings give “net zero” its force: achieving net zero CO<sub>2</sub> emissions, in this sense, is approximately  
147 sufficient to halt CO<sub>2</sub>-induced warming under ambitious mitigation. More complex behaviour<sup>42</sup> may  
148 emerge at much higher levels of warming or much longer timescales.<sup>45</sup>  
149

150 The  $\kappa_E(\rho_F - \rho_E)G\Delta t$  term in equation 1 represents two mutually cancelling processes: a thermal  
151 adjustment ( $\rho_F$ ) and a carbon cycle adjustment ( $\rho_E$ ). If emissions are only reduced to the level  
152 required to stabilise CO<sub>2</sub> concentrations, such that  $E_{\text{GEO}} + E_{\text{LUC}} \approx \rho_E G$  over a multi-decadal period,  
153 then CO<sub>2</sub>-induced warming would continue at a rate  $\rho_F \kappa_E G$ , or about 0.45°C per century if  
154 concentrations are stabilised when temperatures reach 1.5°C (dotted scenario in fig 1 and Extended  
155 Data Fig. 1 a-c). This situation would correspond to all passive CO<sub>2</sub> uptake being included in net zero  
156 calculations. Temperatures would eventually converge to a level determined by the Equilibrium  
157 Climate Sensitivity (ECS),<sup>5,36,37</sup> but the range of uncertainty and especially the risk of a high ECS  
158 remains contested.<sup>36,46-49</sup> Even if atmospheric concentrations were stabilised immediately, the most  
159 likely eventual warming would still exceed 2°C,<sup>50</sup> so simply reducing the net flow of CO<sub>2</sub> into the  
160 atmosphere to zero is not sufficient to limit warming to below 2°C.

161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215

If, however, CO<sub>2</sub> emissions directly resulting from ongoing human activity are reduced to net zero ( $E_{\text{GEO}} + E_{\text{LUC}} = 0$ ) then CO<sub>2</sub>-induced radiative forcing declines at a fractional rate  $\rho_E$  over the following decades (solid scenario in fig 1 and Extended Data Fig. 1 d-f) because of ongoing passive uptake of atmospheric carbon by the oceans and biosphere in response to historical emissions.<sup>12,13</sup> This durable component of passive uptake would continue for many decades even if all human activity were to cease (conversely, if activity continues, measures may be required to protect it). There is no fundamental reason why  $\rho_E = \rho_F$ ,<sup>51</sup> but current best estimates of the difference between them are of order 0.1% per year.<sup>13</sup>

Although the dominant drivers of terrestrial CO<sub>2</sub> uptake are sometimes contested, its overall scale is not. Active net land-use emissions release about 5 GtCO<sub>2</sub> per year into the atmosphere, comprising 7 GtCO<sub>2</sub> per year from deforestation plus 2 GtCO<sub>2</sub> other land cover change minus about 4 GtCO<sub>2</sub> per year due to forest regrowth from past disturbances.<sup>52</sup> In comparison, the current passive land carbon sink is about 12 GtCO<sub>2</sub> per year, estimated from vegetation models, atmospheric inversions, or a simple closure of the global carbon budget.<sup>15,52</sup> How much of this passive land sink is due to CO<sub>2</sub> fertilisation versus other drivers is poorly constrained. The impact of forest demographics, partly an active driver, may be underestimated,<sup>53</sup> which would affect the future of the land sink (demographic changes may saturate sooner than CO<sub>2</sub> fertilisation). Multiple lines of evidence, however, suggest that CO<sub>2</sub> fertilization is likely the single most important driver.<sup>54</sup> When this is added to other passive drivers (temperature and/or precipitation changes, and the passive component of forest regrowth), it becomes likely that the large majority of the global net sink on managed land, as reported in NGHGI and accounted as negative emissions towards countries' emission targets, is passive.

Figure 1 shows a stylized scenario (solid black lines) of global CO<sub>2</sub> emissions,  $E_{\text{GEO}} + E_{\text{LUC}}$ , reduced to net zero in 2050, following the definitions used in those 2009 papers and subsequent IPCC Assessment Reports, hence not including any net passive uptake (solid green lines) in CO<sub>2</sub> removals. This results in CO<sub>2</sub> concentrations peaking before 2050 and declining thereafter, stabilizing global temperatures.<sup>55</sup> Dotted lines show a concentration stabilization scenario in which the net anthropogenic flux of CO<sub>2</sub> into the atmosphere (i.e. the difference between net emissions due to ongoing human activities, dotted grey line in panel a, and net passive uptake in response to historical emissions, or dotted green line) is reduced linearly to zero in 2050 and maintained at zero thereafter. This is sufficient to stabilize atmospheric concentrations but does not halt global warming for many centuries. The dashed lines show a hypothetical "extreme offsetting" scenario in which all passive uptake on land and oceans is progressively re-classified as anthropogenic removals (green shaded area in panel a) and used to offset ongoing emissions to the maximum extent possible to avoid actual emission reductions or active removals. This allows  $E_{\text{GEO}} + E_{\text{LUC}}$  to remain constant past the mid-2030s while nominal emissions, including these offsets, appear to follow the same anthropogenic net-zero pathway as the black solid line. This illustrates the danger of including passive sinks in the definition of net emissions without revisiting climate targets accordingly.<sup>23</sup> Even in the absence of any uncertainty in the climate response, ambiguity in the definition of removals could make the difference between achieving the goals of the Paris Agreement and failing to do so.<sup>24</sup>

[Insert figure 1 here]

If natural systems were to fail to provide the ecosystem service represented by the  $\rho_E G$  term in equation 1, due to Earth system feedbacks or other stresses,<sup>28</sup>  $E_{\text{GEO}} + E_{\text{LUC}}$  would need to be further reduced to  $-\rho_F G$  to prevent further warming. This "equivalent removal" rate is substantial: 0.3% of total historical CO<sub>2</sub> emissions consistent with a peak warming between 1.5 and 2°C (2900-3700 GtCO<sub>2</sub>) is 9-11 GtCO<sub>2</sub> per year.<sup>52</sup> The actual rate of passive CO<sub>2</sub> uptake in the decades after the date of net zero (solid green line in figure 1a) would be about half this equivalent removal rate because active removal of two tonnes of CO<sub>2</sub> is required to reduce the amount of CO<sub>2</sub> in the atmosphere by one tonne.<sup>36</sup> Passive CO<sub>2</sub> uptake plays a bigger role in mitigating the warming impact of ongoing emissions before net zero is achieved, and a smaller role as the carbon cycle begins to re-equilibrate. Yet its continued existence, and the fact that it is not included as a removal in the definition of net

216 anthropogenic emissions, are both essential conditions for net zero CO<sub>2</sub> emissions to halt CO<sub>2</sub>-  
217 induced warming on multi-decadal timescales. Both conditions are potentially at risk.

218

219 **Emerging risks to Net Zero:** The first, unavoidable, risk is that Earth system feedbacks such as  
220 carbon release from thawing permafrost,<sup>57</sup> drying of some wetlands or increased forest fire activity<sup>28,30</sup>  
221 could compromise the net magnitude of biosphere carbon sinks, weakening passive uptake. This effect  
222 is partially accounted for by the use of a constant TCRE, which implies some increase in CO<sub>2</sub>  
223 airborne fraction<sup>20</sup> with cumulative CO<sub>2</sub> emissions cancelling the logarithmic dependence of radiative  
224 forcing on CO<sub>2</sub> concentrations.<sup>42,51,57,58</sup> Even models that represent the full range of Earth system  
225 feedbacks find that this cancellation approximately holds up to 2°C of warming,<sup>59</sup> but it becomes  
226 progressively less certain at higher warming levels<sup>15</sup> and for “overshoot” scenarios.<sup>60</sup> Ultimately, the  
227 only way to minimise the amplifying effect of Earth system feedbacks is to minimise peak warming.  
228 Measures to protect and restore the integrity of biosphere sinks must therefore be additional, not  
229 alternatives, to measures that reduce  $E_{\text{GEO}}$  and  $E_{\text{LUC}}$ . Ongoing fossil fuel emissions and deforestation  
230 put all carbon stored in the biosphere at risk.<sup>61</sup>

231

232 The second “risk” (or moral hazard) arises from policy choices rather than geophysical processes, but  
233 is real nonetheless: unlike the global earth system models and integrated assessment models that  
234 inform IPCC Assessment Reports,<sup>20</sup> greenhouse gas accounting systems, including systems based on  
235 NGHGs<sup>22</sup> and most corporate systems, classify passive uptake that takes place on “managed land”<sup>23</sup>  
236 as an anthropogenic greenhouse gas removal.<sup>62</sup> At present, over 6.5 billion tonnes of CO<sub>2</sub> per year,<sup>62</sup>  
237 or about 60% of total terrestrial carbon uptake,<sup>52</sup> predominantly resulting from passive uptake by  
238 standing forests, are classified as CO<sub>2</sub> removals in national inventories.<sup>23</sup> Most countries define all  
239 their forests as managed for UNFCCC. These accounting systems include this passive uptake in  $E_{\text{LUC}}$ ,  
240 making it available to offset ongoing fossil fuel emissions (Fig. 1, panel a). Indeed, some countries  
241 have used it to declare themselves net zero already.<sup>10</sup>

242

243 These differences in how removals are defined between national inventories and global net zero  
244 pathways are well documented, including by the IPCC.<sup>22–24,62</sup> Although UNFCCC inventory  
245 guidelines<sup>21,63,64</sup> consider all removals on any land declared as managed to be human-induced (i.e.  
246 active), there is potential to add information to NGHGs, including CO<sub>2</sub> uptake on unmanaged land,<sup>65</sup>  
247 that would help countries understand better the magnitude of active and passive components of their  
248 carbon sinks. The availability of this information would make it even more important that the  
249 implications of including passive sinks in emissions targets are understood. It has therefore been  
250 argued<sup>23,24,62</sup> that net emissions in scenarios and targets should be translated to the NGHGI approach  
251 using Dynamic Global Vegetation Models (DGVMs) to include CO<sub>2</sub> uptake on managed lands  
252 explicitly in calculations of  $E_{\text{LUC}}$ , despite inter-DGVM differences.<sup>35</sup> In ambitious mitigation scenarios  
253 the necessary adjustments are small (less than 20%)<sup>23,24</sup> relative to required emission reductions  
254 because only about half to two-thirds of terrestrial carbon uptake is currently classified as taking place  
255 on managed land and passive uptake is expected to decline as emissions fall.<sup>15</sup> Hence, if ambitious  
256 mitigation occurs, ambiguity over passive carbon sinks has an important but limited impact on  
257 allowable emissions at a global level,<sup>23,24</sup> although potentially a much bigger impact at the level of an  
258 individual country or corporation.

259

260 The real problem, however, is that ambiguity in the classification of passive CO<sub>2</sub> uptake may forestall  
261 mitigation getting started. Pressure to classify land as managed (which countries self-determine) will  
262 increase as climate policy requires stronger reductions in net CO<sub>2</sub> emissions. Rising effective carbon  
263 prices increase incentives to monetise all allowable CO<sub>2</sub> removals. The vast majority of countries<sup>62</sup>  
264 already use their managed land sink to assess compliance with emission reduction targets under the  
265 Paris Agreement, even though the Kyoto Protocol attempted to limit<sup>66,67</sup> such use. There is also  
266 increasing interest in monetising “blue carbon” uptake by the oceans.<sup>31</sup> If all passive uptake were  
267 claimed as CO<sub>2</sub> removal, then nominal “net zero CO<sub>2</sub> emissions” would imply only a net zero  
268 atmospheric CO<sub>2</sub> growth rate, or  $E_{\text{GEO}} + E_{\text{LUC}} - \rho_E G = 0$  on multi-decadal timescales. This would  
269 stabilise CO<sub>2</sub> concentrations, which is sufficient to slow further global warming but would not halt it  
270 for centuries. This may seem an extreme scenario (dashed lines in Fig. 1), but it is impossible to

271 predict how accounting conventions will respond to very high effective global carbon prices  
272 associated with ambitious mitigation. A coastal or island state could argue it has a right to take credit  
273 for passive uptake into the oceans of its exclusive economic zone (EEZ) if other countries take credit  
274 for passive uptake into their forests. EEZs account for 30% of global ocean area and an uncertain (but  
275 estimable) fraction of ocean carbon uptake.<sup>68</sup> Credits are already being sold for carbon capture into  
276 the open oceans without clear standards to ensure additionality,<sup>69</sup> raising the prospect of all ocean  
277 passive carbon uptake being claimed as removals, as has already occurred in many regions on land.  
278

279 **How did this situation arise?** Passive CO<sub>2</sub> uptake was not classed as anthropogenic CO<sub>2</sub> removal in  
280 the 2009 papers that established the need for net zero. While the potential role of, and challenge of  
281 quantifying, land-based removals had long been acknowledged,<sup>70</sup> those original papers equated zero  
282 CO<sub>2</sub> emissions with  $E_{\text{GEO}} + E_{\text{LUC}} = 0$  and did not even envisage a substantial negative  $E_{\text{LUC}}$   
283 compensating for ongoing fossil fuel emissions. The only compensatory mechanism considered at that  
284 time for residual fossil use was engineered CO<sub>2</sub> capture (or recapture from the atmosphere) and  
285 geological storage.<sup>71-73</sup>  
286

287 The emphasis on global “net” emissions emerged in the Synthesis Report of the IPCC 5<sup>th</sup> Assessment  
288 (AR5)<sup>74</sup>, but still did not include passive uptake and envisaged a limited role for negative  $E_{\text{LUC}}$ : figure  
289 SPM.14 of that report shows approximately zero net agriculture, forestry and other land-use  
290 (AFOLU) emissions in the majority of technology-neutral mitigation scenarios likely to limit  
291 warming to 2°C. Scenarios limiting warming closer to 1.5°C<sup>75</sup> rely more on negative net AFOLU  
292 emissions but this reliance may be inconsistent with assumed bioenergy use,<sup>76</sup> other sustainable  
293 development goals<sup>77,78</sup> and even international law<sup>79</sup>. This exclusion of passive uptake and limited role  
294 for  $E_{\text{LUC}}$  propagated into the Structured Expert Dialogue (SED)<sup>80</sup> that informed the Paris Agreement.  
295 Annex II, paragraph 69, states: “...if we stop emissions today entirely, there will be no further  
296 warming. Essentially, the commitment to future warming is in future emissions. A stable  
297 concentration, however, will result in further warming.” Crucially, these first two sentences are only  
298 true if passive uptake is not classified as a CO<sub>2</sub> removal, while the final sentence makes clear that  
299 SED participants were aware of the importance of the difference between net zero emissions and net  
300 zero atmospheric CO<sub>2</sub> growth rate.  
301

302 Article 4 of the Paris Agreement<sup>81</sup> does not specify precisely what is included in “removals by sinks”.  
303 While it builds on inventory guidelines used under the UNFCCC and Kyoto Protocol, which treat all  
304 carbon stock changes on managed lands as anthropogenic and hence include some passive uptake in  
305 removals, Article 4 also makes clear that its objective is to deliver Article 2. If “removals” were, in an  
306 extreme case, to include all passive uptake, then achieving the “balance” of Article 4 would imply  
307 only a stabilization of atmospheric CO<sub>2</sub> concentrations (dotted and dashed scenarios in Fig. 1). This  
308 would not halt ongoing warming in time to deliver the goal of Article 2, as was made clear in the  
309 SED. Hence only a restrictive definition of “removals” that excludes passive (indirect) sinks renders  
310 the Paris Agreement’s long-term temperature goal (Art. 2.1a) and the implementing objective (Art.  
311 4.1) jointly consistent with the underlying climate science as it has been understood since 2009.  
312

313 **Scale of the problem:** Figure 2 shows fluxes of CO<sub>2</sub> into and out of the atmosphere under a range of  
314 scenarios. Panel a shows the current situation, with fossil CO<sub>2</sub> emissions and active land-use-change,  
315  $E_{\text{GEO}}$  and  $E_{\text{LUC}}$ , only partially compensated for by passive uptake by land and ocean sinks, leading to a  
316 net accumulation of CO<sub>2</sub> in the atmosphere. All panels illustrate the breakdown of fluxes used in the  
317 2009 papers, in equation 1, and by IPCC Assessment Reports. Under the breakdown used by  
318 NGHGs, 6-7 GtCO<sub>2</sub>/year of the passive land sink in panel a would be reallocated to  $E_{\text{LUC}}$ , reducing it  
319 close to zero.  
320

321 [Insert figure 2 here]  
322

323 Panel b shows the fluxes implied by an instantaneous reduction of fossil fuel emissions by 40-50%  
324 and full compensation of ongoing land-use change emissions with active land-based CO<sub>2</sub> removal.  
325 Atmospheric CO<sub>2</sub> growth rate (pale blue bar) would be reduced to net zero, albeit only momentarily.

326 While the rate of passive uptake would start to decline as soon as CO<sub>2</sub> concentrations stop rising,<sup>56</sup>  
327 this scenario is relevant to net zero claims by sub-global entities, both countries and corporations.  
328 Current accounting rules allow an entity to offset its ongoing emissions against carbon uptake on  
329 managed land, including passive uptake. If all passive uptake were classed as a removal, almost 50%  
330 of global emissions could be fully offset, allowing the entities responsible for them to declare they had  
331 achieved net zero<sup>82</sup> without reducing active emissions at all. If remaining emitters then chose not to  
332 participate in mitigation (plausible, given “ambitious” countries and corporations would be doing  
333 nothing more than offset their emissions against uptake that is occurring anyway), this situation could  
334 persist indefinitely.

336 If the instantaneous balance shown in panel b were achieved globally, passive CO<sub>2</sub> uptake would  
337 decline over the following decades, but emissions would not need to decline all the way to zero to  
338 stabilize atmospheric CO<sub>2</sub> concentrations (panel c, and dotted scenario in fig. 1). Temperatures would  
339 continue to rise at the RACF,  $\rho_F$ . To halt global warming, excess atmospheric CO<sub>2</sub> concentrations  
340 must be allowed to decline by  $\rho_F$ , or 0.3% per year (panel d), corresponding to a total absolute uptake  
341 rate (rate of decrease of atmospheric CO<sub>2</sub> content through both passive uptake and net negative  
342 emissions) of about 5 GtCO<sub>2</sub>/year for peak warming in the range 1.5-2°C.<sup>56</sup> In current Earth System  
343 Models  $\rho_E \approx \rho_F$  so it is sufficient to reduce  $E_{\text{GEO}} + E_{\text{LUC}}$  to net zero to achieve this, but the required  
344 rate of CO<sub>2</sub> decline is set by the need to balance the thermal adjustment, independent of carbon cycle  
345 uncertainties. If current models overstate the scale of passive uptake, then  $E_{\text{GEO}} + E_{\text{LUC}}$  would need to  
346 be net negative to stabilise global temperatures.

348 Over decades, the scope for maintaining a substantial net negative  $E_{\text{LUC}}$  to balance a net positive  $E_{\text{GEO}}$ ,  
349 as in panel d, is limited by earth system feedbacks,<sup>28</sup> the need to balance emissions associated with  
350 food production,<sup>77</sup> and, possibly, the need to compensate for weaker-than-expected passive uptake.  
351 Hence, a durable net zero (panel e and solid scenario in Fig. 1) is likely to require<sup>17</sup> that any remaining  
352 fossil-origin CO<sub>2</sub> production is balanced by CO<sub>2</sub> capture or recapture and geological-timescale  
353 storage, meaning secure storage over multi-century to millennial timescales without ongoing human  
354 intervention. Current evidence suggests that well-managed geological sequestration can meet this  
355 standard.<sup>83</sup> Options such as biochar or biomass burial would need to demonstrate a similar level of  
356 security and durability. So only panel e represents a durable halt to global warming but, if all passive  
357 uptake including blue carbon is treated as an anthropogenic removal, then all four of panels b to e  
358 could be regarded as some kind of net zero CO<sub>2</sub> emissions.

360 **Moving forward:** It is difficult to justify definitions of balance and net zero in individual  
361 commitments that, if replicated globally, would not deliver the Paris Agreement goal of limiting  
362 global warming. Yet<sup>23</sup> it will also be difficult to revise UNFCCC reporting rules to exclude all passive  
363 CO<sub>2</sub> uptake from anthropogenic CO<sub>2</sub> removals. There are genuine issues of capacity, resources and  
364 pragmatism in bringing all countries on board with reporting and accounting following IPCC  
365 Guidelines. Furthermore, many countries are relying on passive uptake to contribute to their emission  
366 goals and may object to its exclusion from international transfers under Article 6 of the Paris  
367 Agreement. Care must also be taken not to jeopardise other benefits of reforestation, such as for  
368 biodiversity.<sup>33</sup> There are, however, some measures that can be taken to mitigate the problem.

370 First, we need wider acknowledgement across both science and policy communities that the problem  
371 exists: achieving and maintaining ‘net zero’ emissions under accounting rules that allow passive CO<sub>2</sub>  
372 uptake to count as CO<sub>2</sub> removal will only slow down global warming. UNFCCC reporting is separate  
373 from target-setting: while countries should be encouraged to report emissions and CO<sub>2</sub> uptake on  
374 managed land, they do not need to treat these “biological” removals as fungible with “geological”  
375 fossil fuel emissions in climate targets.<sup>32</sup> Indeed, accounting methods used by the Kyoto Protocol  
376 discouraged this.<sup>67</sup> Accounting under the Global Stocktake and under Article 6 of the Paris Agreement  
377 should learn from and improve on the Kyoto Protocol approaches to try to separate out what is  
378 “additional” (the result of direct anthropogenic activity) in reported removals.<sup>27</sup> A global effort to  
379 report passive CO<sub>2</sub> uptake separately<sup>65</sup> in greenhouse gas inventories, analogous to separate  
380 specification of short-lived climate pollutants,<sup>84</sup> would help. Discussions have already begun between



381 modellers and inventory compilers on this issue,<sup>62,77</sup> including in the context of the 2024 IPCC Expert  
382 Meeting on Reconciling Land Emissions, and will continue in the 7<sup>th</sup> Assessment Report. At the same  
383 time, countries could be encouraged to document in more detail how passive CO<sub>2</sub> uptake is included  
384 in their approaches to reporting and setting their Nationally Determined Contributions.<sup>24</sup> Such  
385 transparency would allow an assessment of the scale of the problem, and whether it may be increasing  
386 as climate ambition strengthens. It is arguably also in countries' long-term interest to acknowledge the  
387 contribution of passive uptake to their emission goals because, unlike emission reductions or active  
388 removals, passive uptake is contingent on other countries' mitigation decisions: as soon as global CO<sub>2</sub>  
389 emissions start to fall, the rate of uptake in most passive sinks will fall in response.<sup>23</sup>

391 Second, voluntary markets, standard-setters and ambitious countries and corporations can go beyond  
392 the current UNFCCC requirements and exclude passive or indirect uptake from removal credits and  
393 net zero claims. For example, if a source of biomass or an ecosystem is claimed to be carbon neutral,  
394 then the land occupied by that biomass source or ecosystem should absorb CO<sub>2</sub> at the same average  
395 rate that an unmanaged mature ecosystem would absorb CO<sub>2</sub> given current environmental conditions  
396 (location, level and recent rate of increase in atmospheric CO<sub>2</sub> concentrations, climate, etc.). This rate  
397 can either be calculated with a vegetation model or inferred from observations of similar regions: such  
398 methods are already used<sup>26</sup> to assess the extent to which claimed emission reductions are additional to  
399 processes that would have occurred in the absence of an intervention. Even if passive uptake can be  
400 quantified and excluded from claims at an individual project level, however, carbon leakage means  
401 that a clear separation is likely to remain challenging as long as reporting systems are still in  
402 widespread use that allow it to count as a removal.<sup>85</sup>

404 Finally, much of the remaining carbon-absorbing capacity of the biosphere may be required to  
405 compensate for emissions associated with food production, such as fertilizer production and use,  
406 particularly if biological carbon sinks are compromised by climate change itself.<sup>28,86,87</sup> Until it can be  
407 shown that total CO<sub>2</sub> uptake by the biosphere and oceans is large enough to halt CO<sub>2</sub>-induced  
408 warming, it is dangerously optimistic to assume that there will be additional capacity for a negative  
409  $E_{LUC}$  to compensate substantially for ongoing fossil fuel emissions.<sup>13,88</sup> Hence, the third and most  
410 important measure is to recognise the likely long-term infeasibility of balancing substantial ongoing  
411 net positive geological-origin CO<sub>2</sub> emissions with enhanced carbon uptake in the biosphere and  
412 oceans that is genuinely additional to the passive uptake that is already required for net zero emissions  
413 to halt warming. All entities committed to the long-term temperature goal of the Paris Agreement  
414 therefore need to plan to jointly achieve global Geological Net Zero.<sup>13,17,18</sup> This means either  
415 eliminating fossil fuel and fossil carbonate (for cement) use entirely or achieving a balance between  
416 any remaining CO<sub>2</sub> production from geological sources and CO<sub>2</sub> committed to permanent geological  
417 storage, potentially as soon as mid-century. Unlike the biosphere, all significant geological sources  
418 and sinks of CO<sub>2</sub> are unambiguously anthropogenic, clarifying emissions accounting. Acknowledging  
419 the geophysical imperative of Geological Net Zero would allow countries and corporations to future-  
420 proof climate mitigation strategies by planning on a progressive transition to like-for-like balancing of  
421 sources and sinks<sup>17</sup> without waiting for consensus on any change to reporting rules. Differentiating in  
422 greenhouse gas accounting systems between avoided emissions, removals to temporary storage and  
423 removals to permanent storage is, however, essential to track progress to Geological Net Zero.<sup>89</sup>

425 **Responsibility for protection of passive sinks:** Equation 1 also makes clear the paramount  
426 importance of protecting natural CO<sub>2</sub> sinks both during and after the transition to Geological Net  
427 Zero. This will entail opportunity costs, as land or coastal oceans that could be used for food or  
428 bioenergy production are allowed to absorb carbon instead, but this passive uptake cannot be used to  
429 compensate for ongoing fossil fuel emissions if net zero is to achieve a durable halt to global  
430 warming. Fortunately, equation 1 also suggests a possible basis for allocating these costs. To prevent  
431 further warming after emissions reach net zero, annual uptake by passive sinks must be greater than or  
432 equal to  $\phi\rho_F G$ , where  $\phi$  is the Perturbation Airborne Fraction (see Methods).<sup>56</sup> This is approximately  
433 0.15% of cumulative global CO<sub>2</sub> emissions  $G$  over the entire industrial period. Any addition to this  
434 cumulative total increases the size of the passive carbon sink that must be maintained for many  
435 decades after global warming has halted. Whether this causal responsibility translates into a moral or

436 legal responsibility to contribute to maintaining that sink is not a scientific question, but science can  
437 quantify the scale of the challenge: for example, even if the United Kingdom were to achieve net zero  
438 CO<sub>2</sub> emissions before 2050, 0.15% of the U.K.'s contribution to historical cumulative emissions will  
439 be 120 MtCO<sub>2</sub> per year. Should this exceed the passive sink capacity of the U.K.'s land and coastal  
440 oceans,<sup>90</sup> then to genuinely end the U.K.'s contribution to ongoing global warming, the U.K. would  
441 arguably need to undertake active CO<sub>2</sub> removal at approximately double ( $1/\phi$ ) the rate of any  
442 shortfall (in addition to removals to compensate for any ongoing residual emissions) or to rely on  
443 passive uptake in other jurisdictions. Mechanisms for redistributing the costs of maintaining passive  
444 carbon sinks after the date of net zero may therefore be needed.<sup>91</sup> Likewise, undertakings by private  
445 corporations to maintain passive carbon sinks could be seen as addressing the impact of their  
446 historical cumulative emissions, not compensation for future emissions. The traditional concept of  
447 historical responsibility, linking past emissions with future emission reduction rates,<sup>92</sup> remains  
448 complex and multi-faceted.<sup>93</sup> In contrast, the responsibility that we highlight here is a simple  
449 geophysical one: by adding to cumulative emissions, any entity, country or corporation adds to the  
450 total passive carbon sink that needs protection for the foreseeable future.

451  
452 **Actionable implications:** Acknowledging the need for Geological Net Zero makes clear what it takes  
453 for any continued fossil fuel use to be consistent with Paris Agreement goals. Offsetting emissions  
454 with enhanced CO<sub>2</sub> uptake in the oceans and biosphere can provide immediate benefits<sup>33</sup> if and only if  
455 it is genuinely additional to passive CO<sub>2</sub> uptake. In a durable net zero world, 100% of the CO<sub>2</sub>  
456 generated by any continued fossil fuel or fossil carbonate use will almost certainly need to be either  
457 captured at source or recaptured from the atmosphere and committed to geological-timescale storage.  
458 A commitment from high-ambition participants to report and scale up this 'geologically stored  
459 fraction'<sup>94</sup> is needed urgently: it is currently about 0.1% globally,<sup>95</sup> even including CO<sub>2</sub> injection for  
460 enhanced hydrocarbon recovery, and accelerates smoothly over time to reach 100% at the date of  
461 geological net zero in cost-effective scenarios that meet the goals of the Paris Agreement.<sup>96,97</sup> This  
462 implies, in addition to reducing emissions, achieving a 10% geologically stored fraction by the mid  
463 2030s<sup>98</sup> and investing now for a further ten-fold increase in stored fraction over the following 20  
464 years, including demonstrating secure and verifiable geological CO<sub>2</sub> storage capacity to match any  
465 new fossil fuel reserves. These are ambitious but achievable goals for the fossil fuel industry and its  
466 customers.

467 **Figure captions:**

468

469 **Fig 1: Impact of ambiguity in the definition of removals in net zero.** Black and grey lines in panel  
470 a show net CO<sub>2</sub> emissions,  $E_{\text{GEO}} + E_{\text{LUC}}$ , calculated using the definition of removals adopted in IPCC  
471 Assessment Reports (ARs). Green lines show corresponding passive uptake by the oceans and  
472 biosphere. Panels b and c show a central estimate<sup>55</sup> of the response of CO<sub>2</sub> concentrations and global  
473 average surface temperature assuming constant non-CO<sub>2</sub> forcing after 2020 (which requires  
474 immediate rapid reductions in methane emissions to compensate for other changes). Line-styles in all  
475 three panels indicate three scenarios corresponding to different interpretations of net zero. Solid lines  
476 assume net emissions are reduced linearly to zero in 2050, halting warming. Dotted lines assume net  
477 CO<sub>2</sub> flux into the atmosphere (net emissions minus passive uptake) is reduced linearly to zero in  
478 2050, stabilising concentrations. Dashed lines show a scenario that follows the same nominal  
479 emissions pathway as the solid scenario but assumes “reductions” are achieved as far as possible by  
480 reclassifying passive uptake (into both land and oceans) as removals and using it to offset ongoing  
481 (assumed constant) emissions.

482

483 **Fig 2: Fluxes of CO<sub>2</sub> into and out of the atmosphere under different interpretations of net zero.**

484 Red and grey bars indicate energy and industrial emissions and active removal to geological storage,  
485 which net to  $E_{\text{GEO}}$ ; brown and dark green indicate land-use-change emissions and active land-based  
486 removals (using the IPCC Assessment Report definition<sup>20</sup> of removals, including active reforestation  
487 and nature-based solutions), which net to  $E_{\text{LUC}}$ ; light green and dark blue bars indicate passive uptake  
488 by land and oceans; light blue bars indicate net rate of change in the amount of CO<sub>2</sub> in the  
489 atmosphere. (a) present day<sup>52</sup> conditions; (b) fossil fuel emissions reduced instantaneously, but only to  
490 the level required halt the net flow of CO<sub>2</sub> into the atmosphere (mid-21<sup>st</sup>-century dashed scenario in  
491 fig 1); (c) emissions consistent with stable CO<sub>2</sub> concentrations over decades after warming reaches  
492 about 1.5-2°C (dotted scenario in fig 1); (d) emissions consistent with stable temperatures (solid  
493 scenario in fig 1), which requires ongoing passive uptake reducing atmospheric CO<sub>2</sub> (negative pale  
494 blue bar) but allowing some temporary compensation of geological-origin emissions with biogenic  
495 removals; (e) durable net zero, both  $E_{\text{GEO}}$  and  $E_{\text{LUC}}$  equal to zero.

496 **Methods:**

497

498 The origins of equation 1 are detailed in Ref. 18, equations 8 and 14, and summarised here. The total  
 499 anthropogenic change in global average temperature over a multi-decade time-interval is given by the  
 500 following generalisation of equation 1:

501

$$502 \quad \Delta T = \kappa_E [\Delta G + (\rho_F - \rho_E)G\Delta t] + \kappa_F (\Delta F + \rho_F F\Delta t), \quad (2)$$

503

504 where  $\Delta G = (E_{\text{GEO}} + E_{\text{LUC}})\Delta t$  is the total CO<sub>2</sub> emitted or actively removed by human activities over  
 505 the time-interval  $\Delta t$ ,  $G$  is cumulative CO<sub>2</sub> emissions from pre-industrial to around the middle of that  
 506 time-interval,  $\Delta F$  is the change in, and  $F$  is the average, net non-CO<sub>2</sub> radiative forcing, also over that  
 507 time-interval. The Transient Climate Response to Emissions<sup>20</sup> (TCRE)  $\kappa_E = 0.45(\pm 0.18)$  °C per 1,000  
 508 GtCO<sub>2</sub>,<sup>14</sup> while  $\kappa_F = 0.49(\pm 0.1)$  °C per Wm<sup>-2</sup> is the Transient Climate Response to Forcing, or the  
 509 Transient Climate Response<sup>20</sup> (TCR) divided by the radiative forcing due to a doubling of  
 510 atmospheric CO<sub>2</sub> concentrations. The  $\kappa_F \Delta F$  term represents the fast component<sup>36</sup> of the response to  
 511 radiative forcing (defining  $\Delta F$  as the difference between the decade prior to the beginning and the  
 512 decade prior to the end of the time-interval accounts for sub-decadal adjustments), while  $\kappa_F \rho_F F\Delta t$   
 513 represents the gradual adjustment to a constant forcing.<sup>37</sup> Hence the Rate of Adjustment to Constant  
 514 Forcing<sup>18</sup> (RACF)  $\rho_F = (\text{ECS} - \text{TCR})/(\text{TCR} \times s_2)$ , or about 0.3% per year,<sup>40</sup> where ECS is the  
 515 Equilibrium Climate Sensitivity, and  $s_2$  the multi-century adjustment timescale associated with  
 516 warming of the deep oceans<sup>36</sup> and the evolution of feedbacks as the climate system re-equilibrates.<sup>46</sup>

517

518 The  $\kappa_E \Delta G$  term in equation 2 represents the familiar cumulative impact of CO<sub>2</sub> emissions on global  
 519 temperature while the  $\kappa_E (\rho_F - \rho_E)G\Delta t$  term may be understood by considering the limiting case of  
 520  $\rho_E = 0$ : if there were no durable component to passive uptake, and hence CO<sub>2</sub> concentrations and  
 521 CO<sub>2</sub>-induced forcing were to remain constant following net zero emissions, temperatures would  
 522 continue to rise at a fractional rate  $\rho_F$ , or absolute rate  $\kappa_E \rho_F G$ , after an injection of CO<sub>2</sub> taking place  
 523 over a time-scale shorter than  $\rho_F^{-1}$ , which is about 300 years. Studies with coupled climate-carbon-  
 524 cycle models calibrated against available observations<sup>12,13</sup> indicate that temperatures are actually  
 525 expected to change very little after emissions reach net zero: hence  $\rho_E \approx \rho_F$ .

526

527 We now explain the approximations behind the expressions for CO<sub>2</sub>-induced warming in equations 1  
 528 and 2. Over a decade to century time-interval  $\Delta t$  (not longer), the change in atmospheric CO<sub>2</sub> loading  
 529 resulting from anthropogenic CO<sub>2</sub> emissions can be approximated by

530

$$531 \quad \Delta C_A \approx \phi (\Delta G - \rho_E G\Delta t), \quad (3)$$

532

533  $\phi$  being the Perturbation Airborne Fraction, or the change in  $\Delta C_A$  resulting from a unit increase in  $\Delta G$   
 534 over that period.<sup>56</sup> Unlike the instantaneous airborne fraction,  $\Delta C_A/\Delta G$ , which necessarily becomes  
 535 undefined as  $\Delta G \rightarrow 0$ ,  $\phi$  can remain close to its historical value (approximately 50%) even in  
 536 ambitious mitigation scenarios. Similarly, on these timescales, the externally-driven change in global  
 537 mean surface temperature is approximately

538

$$539 \quad \Delta T \approx \kappa_F (\Delta F_{\text{tot}} + \rho_F F_{\text{tot}}\Delta t), \quad (4)$$

540

541 where  $\Delta F_{\text{tot}}$  and  $F_{\text{tot}}$  are, respectively, the change in and average level of total radiative forcing from  
 542 all sources.<sup>36,37</sup> For CO<sub>2</sub>-induced radiative forcing,  $\Delta F_{\text{CO}_2} = \alpha \Delta C_A$ , where  $\alpha$  is the radiative efficacy in  
 543 Wm<sup>-2</sup> per additional billion tonnes of CO<sub>2</sub> in the atmosphere. For emissions concentrated into a time  
 544 much less than  $\rho_E^{-1}$  (as is the case for the historical record), the second term on the right-hand side of  
 545 equation 3 is small, so  $F_{\text{CO}_2} = \alpha \phi G$ . Neither  $\alpha$  nor  $\phi$  is constant, but the non-linearities cancel, such  
 546 that  $\alpha \phi$ , the change in radiative forcing on decade to century timescales per tonne of CO<sub>2</sub> emitted, is  
 547 approximately constant. Substitution of equation 3 into equation 4 and introducing  $\kappa_E = \alpha \phi \kappa_F$  yields  
 548 the expression for CO<sub>2</sub>-induced warming in equations 1 and 2.

549

550 Equation 2 also implies that, before emissions reach net zero, total passive CO<sub>2</sub> uptake by both  
 551 terrestrial biosphere and oceans consists of a transient component (driven by redistribution of recent  
 552 emissions into rapidly-equilibrating carbon reservoirs) and a durable component that is, on multi-  
 553 decade timescales, proportional to cumulative emissions since pre-industrial:<sup>18</sup>

$$554 \quad \Delta G - \Delta C_A \approx [(1 - \phi) \times (E_{\text{GEO}} + E_{\text{LUC}}) + \phi \rho_E G] \Delta t. \quad (5)$$

557 The accuracy of these approximations is illustrated in Extended Data Fig. 1 using the response of the  
 558 FaIR simple climate model<sup>55</sup> to stylized concentration-stabilization and net zero emission scenarios,  
 559 compared with the expressions for passive uptake and temperature response given by equations 5 and  
 560 1, respectively. The FaIR model has been shown<sup>13</sup> to be consistent with the behaviour of much more  
 561 complex Earth System Models over a broad range of scenarios, so agreement with FaIR is indicative  
 562 of agreement with a wider range of models.

564 Under net zero emissions, meaning  $E_{\text{GEO}} + E_{\text{LUC}} = 0$ , the annual rate of passive CO<sub>2</sub> uptake converges  
 565 to  $\phi \rho_E G$ , which has the same impact as active removal of  $\rho_E G$  GtCO<sub>2</sub> per year, or approximately  
 566 0.3% per year of cumulative historical CO<sub>2</sub> emissions. Figure 2 assumes this passive uptake continues  
 567 to be partitioned equally between the terrestrial biosphere and oceans, consistent with the range of  
 568 results of the ZECMIP model intercomparison project (figure 8 of ref. 12). If contributions to the  
 569 protection of these passive sinks were to reflect physical contributions to this committed ongoing  
 570 carbon uptake, research into the geographic location of land and ocean sinks, and the evolution of  
 571 both transient and durable components of passive uptake as emissions decline, is clearly a priority.<sup>90</sup>

573 The level of CO<sub>2</sub>-induced warming after a period of positive emissions starting from pre-industrial  
 574 equilibrium is  $\kappa_E G$  if and only if the time-scale over which those emissions take place is much less  
 575 than  $(\rho_F - \rho_E)^{-1}$ . Since  $\rho_F^{-1} \approx 300$  years and  $\rho_E > 0$ ,  $(\rho_F - \rho_E)^{-1}$  is of order 1,000 years.<sup>18</sup> Hence  
 576 the observation that warming is proportional to cumulative CO<sub>2</sub> emissions for CO<sub>2</sub> injections  
 577 primarily taking place over a century or less (which includes the historical record and most  
 578 experiments used as evidence for this cumulative impact) does not imply that net zero emissions  
 579 would automatically be associated with no further warming or cooling. Likewise, if  $\kappa_E$  is not constant  
 580 (but instead increases with  $G$ , for example), CO<sub>2</sub>-induced warming would still remain constant under  
 581 net zero CO<sub>2</sub> emissions provided  $\rho_F = \rho_E$ . The linear relationship between cumulative CO<sub>2</sub> emissions  
 582 and CO<sub>2</sub>-induced warming is neither necessary nor sufficient for there to be no further warming or  
 583 cooling following net zero CO<sub>2</sub> emissions: these are independent observations, both of which are  
 584 supported by modelling and observations to date.<sup>44</sup>

#### 586 **Extended Data Figure Captions:**

588 Extended Data Fig. 1: **Response to a stylized emission to illustrate the role of passive uptake.** The  
 589 figure shows the response of the FaIR2.0 simple climate model<sup>55</sup> to an emission of 40 billion tonnes  
 590 of CO<sub>2</sub> per year for 70 years, followed by stabilisation of atmospheric concentrations (panels a-c) or  
 591 net zero ongoing emissions (panels d-f). Annual CO<sub>2</sub> flows are shown in panels a and d, changes in  
 592 CO<sub>2</sub> stocks in b and e and temperature response in c and f. Grey, green and blue lines show CO<sub>2</sub>  
 593 emissions, passive uptake and atmospheric increase, annual (panels a and d) and cumulative (panels b  
 594 and e), respectively. Blue and green lines add up to grey lines by construction. Red lines (panels c and  
 595 f) show temperature response. Emissions consistent with stable concentrations are equal to passive  
 596 uptake after concentrations stabilise (panel a) because the rate of atmospheric increase (panel b) is  
 597 then zero. They are initially halved (see fig. 2b of main text), halved again after about 20 years (fig. 2c  
 598 of main text), but do not decline to zero, and temperatures continue to rise for many decades at an  
 599 approximately constant rate (panel c). If emissions are reduced to net zero and passive sinks are not  
 600 compromised, passive uptake immediately draws down the atmospheric CO<sub>2</sub> burden (panels d and e),  
 601 stabilising global temperatures (panel f). Dotted green line shows cumulative passive CO<sub>2</sub> uptake  
 602  $\Delta G - \Delta C_A$  predicted by equation 5 (Methods) with a constant Perturbation Airborne Fraction, PAF,<sup>56</sup>  
 603  $\phi = 0.5$ , and constant Slow Carbon-cycle Adjustment Rate, SCAR,<sup>18</sup>  $\rho_E = 0.3\%$  per year. Dotted red

604 line shows temperature approximated by cumulative emissions, or equation 1 with  $\rho_E = \rho_F$  and  
605 constant Transient Climate Response to Emissions, TCRE,<sup>8</sup>  $\kappa_E$ . These approximations are accurate  
606 relative to the uncertainties in the climate response both while emissions are positive and for the first  
607 few decades after emissions reach net zero, but not over a broader range of timescales and scenarios.  
608

609 **Acknowledgements:** This work was supported by the Strategic Research Fund of the University of  
610 Oxford (MA & SJ), the European Union's Horizon 2020 projects NEGEM (#869192; MA, SJ), 4C  
611 (#821003; MA, PF, GPP), ESM2025 (#101003536; PF, CDJ, JR, RK), PATHFINDER (#101056907;  
612 JH), and PROVIDE (#101003687; JR), the Met Office Hadley Centre Climate Programme funded by  
613 DSIT (JG, CDJ, JAL, PAS), The UKRI programmes GGR-D (NE/V013106/1; JH) and AGILE (CH),  
614 Manchester Metropolitan University (SR), the Research Council of Norway project TRIFECTA  
615 (#334811; GPP), the Swiss National Science Foundation (#200492; TFS), and Environment and  
616 Climate Change Canada's Climate Action and Awareness Fund (NBSClimate, AJW, HDM, KZ). This  
617 paper was initiated through a Fleagle Fellowship in Atmospheric Science Policy at the University of  
618 Washington.  
619

620 **Author contributions:** MA, DF, PF, NG, GG, JG, WH, JH, CH, SJ, CJ, RK, JL, HDM, MM, NM,  
621 GP, GKP, SR, JR, PS, SS, TS, AW and KZ contributed to the drafting and editing of this paper.  
622 Figures were compiled by MA & SJ.  
623

624 **Competing interests declaration:** The authors declare no competing interests. The views expressed  
625 are purely those of the writers and may not under any circumstances be regarded as stating an official  
626 position of the European Commission or any other institution.  
627

628 **Data availability statement:** All data and software required for the reproduction of figures is  
629 provided through CodeOcean [https://codeocean.com/capsule/f7396914-3276-44a6-a7a4-  
630 81df82d2451c/](https://codeocean.com/capsule/f7396914-3276-44a6-a7a4-81df82d2451c/). Datasets include AR6 global radiative forcing timeseries AR6\_ERF\_1750-2019.csv  
631 available on <https://doi.org/10.5285/568fb4b2e6464a50a30c7140bb88a497> and emissions timeseries  
632 Global\_Carbon\_Budget\_2023v1.1.xlsx available on <https://doi.org/10.18160/GCP-2023>  
633

#### 634 **References:**

- 635 1. UNFCCC. Outcome of the first global stocktake. *FCCC/PA/CMA/2023/L.17*  
636 [https://unfccc.int/sites/default/files/resource/cma2023\\_L17\\_adv.pdf?download](https://unfccc.int/sites/default/files/resource/cma2023_L17_adv.pdf?download) (2023).
- 637 2. Parris, H., Anger-Kraavi, A. & Peters, G. P. Does a change in the 'global net zero' language matter? *Global*  
638 *Sustainability* **6**, e13 (2023).
- 639 3. Solomon, S., Plattner, G.-K., Knutti, R. & Friedlingstein, P. Irreversible climate change due to carbon  
640 dioxide emissions. *PNAS* **106**, 1704–1709 (2009).
- 641 **The first of six papers<sup>3–8</sup> published in 2009 recognising the irreversible and cumulative impact of CO<sub>2</sub>**  
642 **emissions on global mean surface temperature and consequent need to reduce CO<sub>2</sub> emissions effectively**  
643 **to zero to limit global warming.**
- 644 4. Meinshausen, M. *et al.* Greenhouse-gas emission targets for limiting global warming to 2°C. *Nature* **458**,  
645 1158–1162 (2009).
- 646 5. Allen, M. R. *et al.* Warming caused by cumulative carbon emissions towards the trillionth tonne. *Nature* **458**,  
647 1163–1166 (2009).
- 648 6. Matthews, H. D., Gillett, N. P., Stott, P. A. & Zickfeld, K. The proportionality of global warming to  
649 cumulative carbon emissions. *Nature* **459**, 829–832 (2009).
- 650 7. Zickfeld, K., Eby, M., Matthews, H. D. & Weaver, A. J. Setting cumulative emissions targets to reduce the  
651 risk of dangerous climate change. *PNAS* **106**, 16129–16134 (2009).
- 652 8. Gregory, J. M., Jones, C. D., Cadule, P. & Friedlingstein, P. Quantifying Carbon Cycle Feedbacks. *Journal of*  
653 *Climate* **22**, 5232–5250 (2009).
- 654 9. Collins, M., Knutti, R. & *et al.* Long-term Climate Change: Projections, Commitments and Irreversibility. in  
655 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment*  
656 *Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, *et al* (eds.)]. 1029–1136*  
657 (Cambridge University Press, 2013).
- 658 10. Hale, T. *et al.* *Net Zero Tracker*. <https://zerotracker.net> (2021).




- 659 11. Masson-Delmotte, V. *Summary for Policymakers. Global Warming of 1.5°C IPCC Special Report on Impacts*  
660 *of Global Warming of 1.5°C above Pre-Industrial Levels in Context of Strengthening Response to Climate*  
661 *Change, Sustainable Development, and Efforts to Eradicate Poverty.* (IPCC, 2018).
- 662 12. MacDougall, A. H., Frölicher, T. L. & Jones, C. D. Is there warming in the pipeline? A multi-model analysis  
663 of the Zero Emissions Commitment from CO<sub>2</sub>. *Biogeosciences* **17**, 2987–3016 (2020).
- 664 **Results of the Zero Emission Commitment Model Intercomparison Project (ZECMIP) demonstrating**  
665 **approximately net zero warming following reduction of CO<sub>2</sub> emissions to zero, further analysed in ref. 13**  
666 **to demonstrate the range of CO<sub>2</sub> emissions consistent with no further warming.**
- 667 13. Jenkins, S. *et al.* The Multi-Decadal Response to Net Zero CO<sub>2</sub> Emissions and Implications for Emissions  
668 Policy. *Geophysical Research Letters* **49**, e2022GL101047 (2022).
- 669 14. Masson-Delmotte, V., P. Zhai, *et al.* (eds.). *Summary for Policymakers. in IPCC AR6 Climate Change 2021,*  
670 *The Physical Science Basis* (Cambridge University Press, 2021).
- 671 15. Canadell, J. G., Monteiro, P. M. S. & *et al.* Chapter 5: Global Carbon and other Biogeochemical Cycles and  
672 Feedbacks. in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the*  
673 *Sixth Assessment Report of the Intergovernmental Panel on Climate Change* 673–816 (Cambridge University  
674 Press, Cambridge, UK, 2021).
- 675 16. Forster, P. M., Storelvmo, T. & *et al.* The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity.  
676 in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth*  
677 *Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai *et al.**  
678 *(eds.)]* (Cambridge University Press, 2021).
- 679 17. Fankhauser, S. *et al.* The meaning of net zero and how to get it right. *Nature Climate Change* **12**, 15–21  
680 (2022).
- 681 **Highlights the importance of different interpretations of net zero, noting that durable net zero requires a**  
682 **like-for-like balance between sources and sinks, with only active removals to permanent geological**  
683 **storage being used to compensate for any ongoing emissions of geological-origin CO<sub>2</sub>, e.g. from burning**  
684 **fossil fuels. See also ref. 19.**
- 685 18. Allen, M. R. *et al.* Net Zero: Science, Origins, and Implications. *Annual Review of Environment and*  
686 *Resources* **47**, 849–887 (2022).
- 687 **Reviews the science of net zero, demonstrating the central role of the compensation between the Rate of**  
688 **Adjustment to Constant Forcing and Slow Carbon-cycle Adjustment Rate, or rate of CO<sub>2</sub> forcing decline**  
689 **under zero emissions, and introducing the conceptual framework of equation 1.**
- 690 19. Rogelj, J. Net zero targets in science and policy. *Environ. Res. Lett.* **18**, 021003 (2023).
- 691 20. Matthews, J. B. R., Möller, V., van Diemen, R. & *et al.* IPCC, 2021: Annex VII: Glossary. in *Climate Change*  
692 *2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the*  
693 *Intergovernmental Panel on Climate Change e [Masson-Delmotte, V., P. Zhai, *et al.* (eds.)].* 2215–2256  
694 (Cambridge University Press, Cambridge, UK, and New York, USA, 2021).
- 695 21. Eggleston, S., Buendia, L., Miwa, K. & *et al.* (eds). *2006 IPCC Guidelines for National Greenhouse Gas*  
696 *Inventories Volume 4: Agriculture, Forestry and Other Land Use.* (IGES, Japan, 2006).
- 697 22. Grassi, G. *et al.* Reconciling global-model estimates and country reporting of anthropogenic forest CO<sub>2</sub>  
698 sinks. *Nature Climate Change* **8**, 914–920 (2018).
- 699 23. Grassi, G., Stehfest, E., Rogelj, J. & van Vuuren, D. Critical adjustment of land mitigation pathways for  
700 assessing countries’ climate progress. *Nature Climate Change* **11**, 425–434 (2021).
- 701 **Details correction required to global Paris-aligned pathways to account for national reporting of passive**  
702 **uptake on managed land as a CO<sub>2</sub> removal, developing the issues raised by ref. 22 and further updated in**  
703 **refs. 24 and 62.**
- 704 24. Gidden, M. J. *et al.* Aligning climate scenarios to emissions inventories shifts global benchmarks. *Nature*  
705 **624**, 102–108 (2023).
- 706 25. Canadell, J. G. *et al.* Factoring out natural and indirect human effects on terrestrial carbon sources and sinks.  
707 *Environmental Science & Policy* **10**, 370–384 (2007).
- 708 26. West, T. A. P., Börner, J., Sills, E. O. & Kontoleon, A. Overstated carbon emission reductions from voluntary  
709 REDD+ projects in the Brazilian Amazon. *Proceedings of the National Academy of Sciences* **117**, 24188–  
710 24194 (2020).
- 711 27. Smith, S. M. *et al.* *The State of Carbon Dioxide Removal - 2nd Edition.* <https://osf.io/f85qj/> (2024).
- 712 28. Zickfeld, K. *et al.* Net-zero approaches must consider Earth system impacts to achieve climate goals. *Nat.*  
713 *Clim. Chang.* **13**, 1298–1305 (2023).
- 714 **Highlights the risk that Earth system feedbacks weaken terrestrial carbon sinks, limiting the degree to**  
715 **which climate goals can be met through balancing ongoing emissions with terrestrial uptake.**
- 716 29. Ke, P., Ciais, P., Sitch, S. & *et al.* Low latency carbon budget analysis reveals a large decline of the land  
717 carbon sink in 2023. *Nature* **634**, (2024).


- 718 30. Ke, P. *et al.* Low latency carbon budget analysis reveals a large decline of the land carbon sink in 2023.  
719 *National Science Review* nwae367 (2024) doi:10.1093/nsr/nwae367.
- 720 31. Bertram, C. *et al.* The blue carbon wealth of nations. *Nat. Clim. Chang.* **11**, 704–709 (2021).
- 721 32. den Elzen, M. G. J. *et al.* Updated nationally determined contributions collectively raise ambition levels but  
722 need strengthening further to keep Paris goals within reach. *Mitig Adapt Strateg Glob Change* **27**, 33 (2022).
- 723 33. Seddon, N. *et al.* Understanding the value and limits of nature-based solutions to climate change and other  
724 global challenges. *Philosophical Transactions of the Royal Society B: Biological Sciences* **375**, 20190120  
725 (2020).
- 726 34. Girardin, C. A. J. *et al.* Nature-based solutions can help cool the planet — if we act now. *Nature* **593**, 191–  
727 194 (2021).
- 728 35. Ruehr, S. *et al.* Evidence and attribution of the enhanced land carbon sink. *Nat Rev Earth Environ* **4**, 518–  
729 534 (2023).
- 730 36. Held, I. M. *et al.* Probing the Fast and Slow Components of Global Warming by Returning Abruptly to  
731 Preindustrial Forcing. *Journal of Climate* **23**, 2418–2427 (2010).
- 732 37. Seshadri, A. K. Fast–slow climate dynamics and peak global warming. *Clim Dyn* **48**, 2235–2253 (2017).
- 733 38. Seshadri, A. K. Origin of path independence between cumulative CO<sub>2</sub> emissions and global warming. *Clim*  
734 *Dyn* **49**, 3383–3401 (2017).
- 735 39. Allen, M. R. & Dube, O. P. Chapter 1, Framing and Context. in *Global Warming of 1.5°C. An IPCC Special*  
736 *Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse*  
737 *gas emission pathways [Masson-Delmotte, V., Zhai, P., et al (eds.)]* 49–92 (Cambridge University Press,  
738 Cambridge, UK, and New York, USA, 2018).
- 739 40. Cain, M. *et al.* Improved calculation of warming-equivalent emissions for short-lived climate pollutants. *npj*  
740 *Climate and Atmospheric Science* **2**, (2019).
- 741 41. Matthews, H. D., Solomon, S. & Pierrehumbert, R. Cumulative carbon as a policy framework for achieving  
742 climate stabilization. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and*  
743 *Engineering Sciences* **370**, 4365–4379 (2012).
- 744 42. Herrington, T. & Zickfeld, K. Path independence of climate and carbon cycle response over a broad range of  
745 cumulative carbon emissions. *Earth System Dynamics* **5**, 409–422 (2014).
- 746 43. MacDougall, A. H. & Friedlingstein, P. The Origin and Limits of the Near Proportionality between Climate  
747 Warming and Cumulative CO<sub>2</sub> Emissions. *Journal of Climate* **28**, 4217–4230 (2015).
- 748 44. Seshadri, A. K. Cumulative emissions accounting of greenhouse gases due to path independence for a  
749 sufficiently rapid emissions cycle. *Clim Dyn* **57**, 787–798 (2021).
- 750 45. Frölicher, T. L., Winton, M. & Sarmiento, J. L. Continued global warming after CO<sub>2</sub> emissions stoppage.  
751 *Nature Clim Change* **4**, 40–44 (2014).
- 752 46. Armour, K. C., Bitz, C. M. & Roe, G. H. Time-Varying Climate Sensitivity from Regional Feedbacks.  
753 *Journal of Climate* **26**, 4518–4534 (2013).
- 754 47. Sherwood, S. C. *et al.* An Assessment of Earth’s Climate Sensitivity Using Multiple Lines of Evidence.  
755 *Reviews of Geophysics* **58**, e2019RG000678 (2020).
- 756 48. Huntingford, C., Williamson, M. S. & Nijssen, F. J. M. M. CMIP6 climate models imply high committed  
757 warming. *Climatic Change* **162**, 1515–1520 (2020).
- 758 49. Lewis, N. Objectively combining climate sensitivity evidence. *Clim Dyn* **60**, 3139–3165 (2023).
- 759 50. Abrams, J. F. *et al.* Committed Global Warming Risks Triggering Multiple Climate Tipping Points. *Earth’s*  
760 *Future* **11**, e2022EF003250 (2023).
- 761 51. Williams, R. G., Goodwin, P., Rousenov, V. M. & Bopp, L. A framework to understand the transient climate  
762 response to emissions. *Environ. Res. Lett.* **11**, 015003 (2016).
- 763 **The first complete conceptual framework to understand the cumulative impact of CO<sub>2</sub> emissions on**  
764 **global temperatures and the balance between thermal and carbon cycle adjustments that results in no**  
765 **further CO<sub>2</sub>-induced warming following net zero CO<sub>2</sub> emissions.**
- 766 52. Friedlingstein, P. *et al.* Global Carbon Budget 2023. *Earth System Science Data* **15**, 5301–5369 (2023).
- 767 53. Yang, H. *et al.* Global increase in biomass carbon stock dominated by growth of northern young forests over  
768 past decade. *Nat. Geosci.* **16**, 886–892 (2023).
- 769 54. Walker, A. P. *et al.* Integrating the evidence for a terrestrial carbon sink caused by increasing atmospheric  
770 CO<sub>2</sub>. *New Phytologist* **229**, 2413–2445 (2021).
- 771 55. Leach, N. J. *et al.* FaIRv2.0.0: A generalized impulse response model for climate uncertainty and future  
772 scenario exploration. *Geoscientific Model Development* **14**, 3007–3036 (2021).
- 773 56. Jones, C. D. *et al.* Simulating the Earth system response to negative emissions. *Environ. Res. Lett.* **11**,  
774 095012 (2016).
- 775 57. MacDougall, A. H. Estimated effect of the permafrost carbon feedback on the zero emissions commitment to  
776 climate change. *Biogeosciences* **18**, 4937–4952 (2021).




- 777 58. Millar, R., Allen, M., Rogelj, J. & Friedlingstein, P. The cumulative carbon budget and its implications.  
778 *Oxford Review of Economic Policy* **32**, 323–342 (2016).
- 779 59. Williams, R. G., Ceppi, P. & Katavouta, A. Controls of the transient climate response to emissions by  
780 physical feedbacks, heat uptake and carbon cycling. *Environ. Res. Lett.* **15**, 0940c1 (2020).
- 781 60. Tachiiri, K., Hajima, T. & Kawamiya, M. Increase of the transient climate response to cumulative carbon  
782 emissions with decreasing CO<sub>2</sub> concentration scenarios. *Environ. Res. Lett.* **14**, 124067 (2019).
- 783 61. Hubau, W. *et al.* Asynchronous carbon sink saturation in African and Amazonian tropical forests. *Nature* **579**,  
784 80–87 (2020).
- 785 62. Grassi, G. *et al.* Harmonising the land-use flux estimates of global models and national inventories for 2000–  
786 2020. *Earth System Science Data* **15**, 1093–1114 (2023).
- 787 63. IPCC. *Revisiting the Use of Managed Land as a Proxy for Estimating National Anthropogenic Emissions and*  
788 *Removals – IPCC Expert Meeting Report.* (2010).
- 789 64. IPCC. *Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories.* (2019).
- 790 65. Nabuurs, G.-J., Ciais, P., Grassi, G., Houghton, R. A. & Sohngen, B. Reporting carbon fluxes from  
791 unmanaged forest. *Commun Earth Environ* **4**, 1–4 (2023).
- 792 66. Fry, I. Twists and Turns in the Jungle: Exploring the Evolution of Land Use, Land-Use Change and Forestry  
793 Decisions within the Kyoto Protocol. *Review of European Community & International Environmental Law*  
794 **11**, 159–168 (2002).
- 795 67. Macintosh, A. K. LULUCF in the post-2012 regime: fixing the problems of the past? *Climate Policy* **12**,  
796 341–355 (2012).
- 797 68. Gruber, N. *et al.* Trends and variability in the ocean carbon sink. *Nat Rev Earth Environ* **4**, 119–134 (2023).
- 798 69. Reimers, J. B. P., Jessica Cross, Matthew C. Long, Patrick A. Rafter, Clare E. The Science We Need to  
799 Assess Marine Carbon Dioxide Removal. *Eos* [http://eos.org/opinions/the-science-we-need-to-assess-marine-](http://eos.org/opinions/the-science-we-need-to-assess-marine-carbon-dioxide-removal)  
800 [carbon-dioxide-removal](http://eos.org/opinions/the-science-we-need-to-assess-marine-carbon-dioxide-removal) (2023).
- 801 70. Prentice, I. C. & *et al.* Chapter 3: The Carbon Cycle and Atmospheric Carbon Dioxide. in *Climate Change*  
802 *2001: The Scientific Basis, The IPCC Third Assessment Report [Houghton, J.T., Ding, Y. *et al* (eds)]* 185–  
803 237 (Cambridge University Press, 2001).
- 804 71. Weaver, A. J., Zickfeld, K., Montenegro, A. & Eby, M. Long term climate implications of 2050 emission  
805 reduction targets. *Geophysical Research Letters* **34**, (2007).
- 806 72. Allen, M. *et al.* The exit strategy. *Nature Clim Change* **1**, 56–58 (2009).
- 807 73. Allen, M. R., Frame, D. J. & Mason, C. F. The case for mandatory sequestration. *Nature Geoscience* **2**, 813–  
808 814 (2009).
- 809 74. Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.). *IPCC, 2014: Climate Change 2014: Synthesis*  
810 *Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental*  
811 *Panel on Climate Change.* 151 pp (2014).
- 812 75. Skea, J., Shukla, P. R. & *al.* Summary for Policymakers. in *Climate Change 2022: Mitigation of Climate*  
813 *Change. Contribution of Working Group III to the IPCC Sixth Assessment Report (AR6)* (Cambridge  
814 University Press, 2022).
- 815 76. Harper, A. B. *et al.* Land-use emissions play a critical role in land-based mitigation for Paris climate targets.  
816 *Nat Commun* **9**, 2938 (2018).
- 817 **Identifies the essential role played by the land carbon sink in 1.5°C scenarios and raises questions about**  
818 **compatibility with other sustainable development goals.**
- 819 77. Shukla, P. R., Skea, J. & Calvo Buendia, E. *Climate Change and Land: An IPCC Special Report on Climate*  
820 *Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse*  
821 *Gas Fluxes in Terrestrial Ecosystems.* (IPCC, 2019).
- 822 78. Cobo, S., Galán-Martín, Á., Tulus, V., Huijbregts, M. A. J. & Guillén-Gosálbez, G. Human and planetary  
823 health implications of negative emissions technologies. *Nat Commun* **13**, 2535 (2022).
- 824 **Documents the potential conflict between human and ecosystem health and heavy dependence on**  
825 **terrestrial carbon removal to achieve climate goals.**
- 826 79. Stuart-Smith, R. F., Rajamani, L., Rogelj, J. & Wetzer, T. Legal limits to the use of CO<sub>2</sub> removal. *Science*  
827 **382**, 772–774 (2023).
- 828 80. UNFCCC. *Report on the Structured Expert Dialogue on the 2013–2015 Review. Note by the Co-Facilitators*  
829 *of the Structured Expert Dialogue.* <https://unfccc.int/documents/8707> (2015).
- 830 **Report of the 2013-14 Structured Expert Dialogue providing essential context of how net zero science was**  
831 **communicated to the negotiators of the Paris Agreement.**
- 832 81. UNFCCC. *Adoption of the Paris Agreement.* 27  
833 [https://unfccc.int/sites/default/files/english\\_paris\\_agreement.pdf](https://unfccc.int/sites/default/files/english_paris_agreement.pdf) (2015).
- 834 82. Mitchell-Larson, E. & Bushman, T. *Carbon Direct Commentary: Release of the Voluntary Registry Offsets*  
835 *Database.* [https://carbon-direct.com/wp-content/uploads/2021/04/CD-Commentary-on-Voluntary-Registry-](https://carbon-direct.com/wp-content/uploads/2021/04/CD-Commentary-on-Voluntary-Registry-Offsets-Database_April-2021.pdf)  
836 [Offsets-Database\\_April-2021.pdf](https://carbon-direct.com/wp-content/uploads/2021/04/CD-Commentary-on-Voluntary-Registry-Offsets-Database_April-2021.pdf) (2021).

- 837 83. Daniels, S. & et al. *Deep Geological Storage of CO<sub>2</sub> on the UK Continental Shelf: Containment Certainty*.  
838 <https://www.gov.uk/government/publications/deep-geological-storage-of-carbon-dioxide-co2-offshore-uk->  
839 [containment-certainty](https://www.gov.uk/government/publications/deep-geological-storage-of-carbon-dioxide-co2-offshore-uk-) (2023).
- 840 84. Allen, M. R. *et al.* Indicate separate contributions of long-lived and short-lived greenhouse gases in emission  
841 targets. *npj Clim Atmos Sci* **5**, 1–4 (2022).
- 842 85. Searchinger, T. D., Wiersenius, S., Beringer, T. & Dumas, P. Assessing the efficiency of changes in land use  
843 for mitigating climate change. *Nature* **564**, 249–253 (2018).
- 844 86. Skea, J. *IPCC Summary for Policymakers*. In *Climate Change 2022: Mitigation of Climate Change*.  
845 *Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on*  
846 *Climate Change*. 1–50  
847 [https://www.ipcc.ch/report/ar6/wg3/downloads/report/IPCC\\_AR6\\_WGIII\\_SummaryForPolicymakers.pdf](https://www.ipcc.ch/report/ar6/wg3/downloads/report/IPCC_AR6_WGIII_SummaryForPolicymakers.pdf)  
848 (2022).
- 849 87. Nabuurs, G.-J., Mrabet, R. & et al. Chapter 7: Agriculture, Forestry, and Other Land Uses (AFOLU). in  
850 *Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth*  
851 *Assessment Report of the Intergovernmental Panel on Climate Change*[P.R. Shukla, J. Skea, et al (eds.)]  
852 747–860 (Cambridge University Press, Cambridge, UK, and New York, USA, 2022).
- 853 88. Matthews, H. D. *et al.* Temporary nature-based carbon removal can lower peak warming in a well-below 2  
854 °C scenario. *Commun Earth Environ* **3**, 1–8 (2022).
- 855 89. Schenuit, F. *et al.* Secure robust carbon dioxide removal policy through credible certification. *Commun Earth*  
856 *Environ* **4**, 1–4 (2023).
- 857 90. Chandra, N. *et al.* Estimated regional CO<sub>2</sub> flux and uncertainty based on an ensemble of atmospheric CO<sub>2</sub>  
858 inversions. *Atmospheric Chemistry and Physics* **22**, 9215–9243 (2022).
- 859 91. Ciais, P. *et al.* Attributing the increase in atmospheric CO<sub>2</sub> to emitters and absorbers. *Nature Clim Change* **3**,  
860 926–930 (2013).
- 861 92. Shue, H. Historical Responsibility, Harm Prohibition, and Preservation Requirement: Core Practical  
862 Convergence on Climate Change. *Moral Philosophy and Politics* **2**, 7–31 (2015).
- 863 93. Sardo, M. C. Responsibility for climate justice: Political not moral. *European Journal of Political Theory* **22**,  
864 26–50 (2023).
- 865 94. Jenkins, S., Mitchell-Larson, E., Ives, M. C., Haszeldine, S. & Allen, M. Upstream decarbonization through a  
866 carbon takeback obligation: An affordable backstop climate policy. *Joule* **5**, 2777–2796 (2021).
- 867 95. Steyn, M., Oglesby, J., Turan, G. & et al. *Global Status of CCS 2022*.  
868 [https://status22.globalccsinstitute.com/wp-content/uploads/2023/03/GCCSI\\_Global-Report-](https://status22.globalccsinstitute.com/wp-content/uploads/2023/03/GCCSI_Global-Report-2022_PDF_FINAL-01-03-23.pdf)  
869 [2022\\_PDF\\_FINAL-01-03-23.pdf](https://status22.globalccsinstitute.com/wp-content/uploads/2023/03/GCCSI_Global-Report-2022_PDF_FINAL-01-03-23.pdf) (2023).
- 870 96. Jenkins, S., Mitchell-Larson, E., Ives, M. C., Haszeldine, S. & Allen, M. Upstream decarbonization through a  
871 carbon takeback obligation: An affordable backstop climate policy. *Joule* **5**, 2777–2796 (2021).
- 872 97. Jenkins, S., Kuijper, M., Helferty, H., Girardin, C. & Allen, M. Extended producer responsibility for fossil  
873 fuels. *Environ. Res. Lett.* **18**, 011005 (2023).
- 874 98. Skidmore, C. *Mission Zero - Independent Review of Net Zero*. 339  
875 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/1128689/m](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1128689/m)  
876 [ission-zero-independent-review.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1128689/m) (2022).
- 877







Extended Data Fig. 1